

# POINTER™ ALB INSECTICIDE

## Specimen Label

For Use Only Under U.S.D.A. Supervision with U.S.D.A.-Approved Injection Devices

### ACTIVE INGREDIENT:

Imidacloprid (CAS No. 138261-41-3) ..... 5%

OTHER INGREDIENTS ..... 95%

Total: 100%

Container Size	Net Contents	AI/Container
1000 ml pack	1 qt 2 fl oz (1000 ml)	50 grams

EPA Reg. No. 69117-1 EPA Est. 69117-NE-1

Keep Out of Reach of Children

## WARNING

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals:

**WARNING:** Harmful if swallowed, inhaled or absorbed through skin. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear safety glasses. Thoroughly wash with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### FIRST AID

**If Swallowed:** Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**Have the product container or label with you when calling a poison control center or doctor, or going for treatment.**

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

For assistance contact



P.O. Box 34645 • Omaha, NE 68134 • www.ArborSystems.com  
Toll-Free: 1-800-698-4641 • Phone: (402) 339-4459 • FAX: (402) 339-5011

#### NOTICE OF WARRANTY

ArborSystems warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. ArborSystems does not make or authorize any agent or representative to make any other warranty, guarantee or representation, express or implied, concerning this product.

Pointer™ Insecticide is a trademark of ArborSystems.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its label.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Use protective eyewear and gloves when handling this product. Use chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

#### U.S.D.A. RATE SPECIFICATIONS FOR ASIAN AND CITRUS LONGHORNED BEETLE PROGRAMS. For Use Only Under U.S.D.A. Supervision. For Use with U.S.D.A.-Approved Injection Devices

Read and follow the manufacturers' written documentation on the proper use and maintenance of the injector you are using. The following rates are generally recommended as a function of tree diameter at breast height (DBH).

2 to 11 inches DBH – 4 ml per diameter inch.

12 to 23 inches DBH – 8 ml per diameter inch.

24 to 35 inches DBH – 16 ml per diameter inch.

36 inches and above DBH – 24 ml per diameter inch.

#### GENERAL DIRECTIONS

To determine the initial number of needed injection sites, measure the tree at breast height using a standard forestry tape measure.

- If measuring the circumference, divide this number by 6 to determine the number of sites needed.
- If measuring the diameter, divide this number by 2 to determine the number of injection sites needed.
- Initial injector sites should be in the active sapwood, evenly spaced around the tree at the root buttress region and avoiding the root valleys.
- If a single site cannot accept its full dose, that dose may be divided among the other initial injection sites.
- If additional injection sites are needed they may be spaced horizontally or vertically from the initial sites.

The minimum horizontal spacing for injection sites is 3 inches. The minimum vertical spacing for injection sites is 6 inches. Preferably vertical spacing should be staggered and not aligned. Do not apply more than once per calendar year.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store above 65°F in a cool, dry place and in such a manner as to prevent cross contamination with fertilizers, food, feed and other pesticides. Store in original container out of reach of children, preferably in a locked storage area.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration; or if allowed by local authorities, by burning; if burned stay out of smoke.